



Smart Solutions for Today's Geoscientist

BLOCK: BIN QASIM SOUTH

OFFSHORE BIDDING BLOCK ROUND 2023


MINISTRY OF ENERGY PETROLEUM DIVISION (DGPC)

www.ppisonline.com

INTRODUCTION



- Bin Qasim South Block covers an area of 2021.69 Sq. Kms.
- Geological Basin: Offshore Indus, Basin Pakistan.
- The block falls in Prospectivity Zone O.
- Lasmo, Husky, PC, Wintershall, OXY and OGDCL acquired 2D and 3D seismic data approximately 8215.5424 L. Kms and 339.90 Sq. Kms in the block within the years 1969, 1972, 1976, 1977, 1982, 1986, 1989, 1998 and 2009 respectively.
- The Block is surrounded by Gharo Creek Block in the South.
- The wells drilled in the near vicinity is Indus Marine 1B, Shark 01, Dabbo Creek 01 and Patiani Creek 01.





GEOLOGICAL HISTORY



Late Cretaceous – Early Paleocene:

- Rapid northward movement of Indian Plate after separation from Madagascar
- Bela ophiolites obduction
- Extrusion of Deccan Volcanics

Paleocene - Eocene:

 Deposition of limestone on seamounts and shales in lows / depressions


Miocene

Oligocene

Oligo-Miocene:

Present Day

- Himalayan orogeny
- Indus Delta-Fan deposition





PETROLEUM SYSTEM



Source Rock:

- Lower Cretaceous Sembar can be the source rock.
- Paleocene carbonates and Miocene shales can act as source rock.

Reservoir Rock:


 Upper Cretaceous Pab and Paleocene Ranikot Formations can be reservoir targets in near shore areas.

Seal:

Mud dominated sediments and intraformational shales packages will provide seals for potential reservoirs.

Trap:

- Both structural and stratigraphic traps
- Eocene carbonates build ups over seamounts, growth faults, roll-over anticlines and stratigraphic traps within deltaic fan system (e.g., bars, barrier islands and pinch-out / facies change etc.) are likely trapping mechanisms







Syed Asif Ahsan et,al., (2012) Physico-Chemical Controls on Source Rock in Offshore Indus – Comparative Study of Some Major Tertiary Deltas of the World : PAPG/SPE Annual Technical Conference WWW.ppisonline.com

PROSPECTIVITY





- High resolution seismic data can allow to delineate true potential of the block
- Both structural and stratigraphic traps





EXPLORATION RISKS



- Source & Charge: Medium to High risk
- Reservoir: Low to Medium risk
- Seal: Low to Medium risk
- Trap: Low to Medium risk
- Key challenges for future exploration in Tertiary Petroleum System are to establish:
 - Distribution and timing of effective source intervals' development within the drainage area of prospect.
 - Timing of over-pressuring (up to 7000 psi at 2800m in Indus Marine-1A well) within Miocene section (for Miocene and younger targets) with respect to source rock maturation and expulsion.





Comparison suggests that discoveries in offshore deltas have been made in:

- Extension of proven onshore petroleum system to offshore at drillable depth (e.g. Niger, Nile, Irrawady & Mahakam deltas)
- Reservoir –Seal pairs associated with good quality, but less mature source rock drilled onshore (at shallow depths) progressively mature in offshore (e.g. Krishna-Godavri and Nile deltas)
- Biogenic gas found in shallow younger Tertiary section (e.g. Krishna-Godavri and Nile deltas)

International offshore exploration efforts in deltaic areas have generally been successful due to:

- Extension of established onshore petroleum system to offshore at drillable depths
- Good quality less mature source rock drilled onshore progressively mature in offshore
- Gas discoveries of biogenic origin



OFFSHORE BIDDING BLOCK ROUND 2023

MINISTRY OF ENERGY PETROLEUM DIVISION (DGPC)

